

(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 927 717 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:
07.07.1999 Bulletin 1999/27

(51) Int. Cl.⁶: C07C 409/10, C07C 409/08,
C07C 409/00, C07C 407/00

(21) Application number: 98929761.9

(86) International application number:
PCT/JP98/02935

(22) Date of filing: 30.06.1998

(87) International publication number:
WO 99/02491 (21.01.1999 Gazette 1999/03)

(84) Designated Contracting States:
BE DE ES FR GB IT NL

(30) Priority: 09.07.1997 JP 18329197

(71) Applicant:
Mitsui Chemicals, Inc.
Tokyo 100-6070 (JP)

(72) Inventors:
• MATSUI, Shigekazu
Kuga-gun, Yamaguchi-ken 740-0061 (JP)
• KURODA, Hiroshi
Kuga-gun, Yamaguchi-ken 740-0061 (JP)
• HIROKANE, Nobuya
Kuga-gun, Yamaguchi-ken 740-0061 (JP)

• MAKIO, Haruyuki
Kuga-gun, Yamaguchi-ken 740-0061 (JP)
• TAKAI, Toshihiro
Kuga-gun, Yamaguchi-ken 740-0061 (JP)
• KATO, Koji
Kuga-gun, Yamaguchi-ken 740-0061 (JP)
• FUJITA, Terunori
Kuga-gun, Yamaguchi-ken 740-0061 (JP)
• KAMIMURA, Makoto
Kuga-gun, Yamaguchi-ken 740-0061 (JP)

(74) Representative: HOFFMANN - EITLE
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(54) PROCESS FOR PRODUCING HYDROPEROXIDES

(57) A process for preparing hydroperoxides which comprises oxidizing hydrocarbon by a gas containing oxygen in the presence of a specific compound and converting them selectively to corresponding hydroperoxides.

The specific compound is the compound that can capture radicals. The preferable example may be a compound selected from radicals of oxygen, nitrogen, phosphorus, sulfur, carbon or silicon or a compound that forms radicals of these in the reaction system.

The present invention can be applied to oxidation of hydrocarbons including arylalkylhydrocarbons such as cumene, m-diisopropylbenzene, p-diisopropylbenzene, 1,3,5-triisopropylbenzene, isopropyl-naphthalene, diisopropyl-naphthalene, isopropylbiphenyl, diisopropylbiphenyl, etc.

EP 0 927 717 A1

Description

TECHNICAL FIELD

- 5 [0001] The present invention relates to a process for preparing hydroperoxides, which comprises oxidizing hydrocarbon by a gas containing oxygen and converting it selectively to corresponding hydroperoxides.

TECHNICAL BACKGROUND

- 10 [0002] The process of oxidizing hydrocarbon by a gas containing oxygen in the absence of a catalyst to prepare corresponding hydroperoxides is known as autoxidation technology. However, in this reaction, it is necessary to raise the reaction temperature to increase the rate of accumulation of hydroperoxides. Nevertheless, when the reaction temperature is raised to increase the rate of accumulation, the reaction product hydroperoxide is thermally decomposed, resulting in a decline in its selectivity. In other words, since there was such relationship between the rate of accumulation and selectivity that as either one goes up, the other goes down, it was difficult to maintain both of them at high levels.
- [0003] Attempts have been made to oxidize hydrocarbon in the state of gas containing oxygen by a catalyst to produce an effect on the rate of accumulation and/or selectivity in producing corresponding hydroperoxides (Japanese Patent Publication SHO 55-50020, for example).
- 20 [0004] The inventors of the present invention firmly believed that there should be a way to achieve high selectivity, while overcoming the relationship between the rate of accumulation and selectivity and maintaining a commercially viable rate of accumulation at the same time. With this belief, the inventors continued research assiduously. As a result, the inventors successfully made an invention to solve the problem.
- [0005] In the course to the invention, inventors found that a oxidation of a hydrocarbon by a gas containing oxygen in the presence of a specific compound is effective for converting it to the corresponding hydroperoxides.
- 25 [0006] Specifically, the present invention provides a method for oxidizing hydrocarbon by a gas containing oxygen thereby to prepare corresponding hydroperoxides at a high selectivity.

DISCLOSURE OF THE INVENTION


- 30 [0007] The process for preparing hydroperoxides by the present invention is based on the attainment of a very high selectivity possible by the use of a compound capable of capturing radicals as a specified compound in the process of oxidizing hydrocarbon by a gas containing oxygen. The present invention is characterized by oxidizing hydrocarbon by a gas containing oxygen in the presence of a compound that can capture radicals and converting it selectively to corresponding hydroperoxides.
- 35

THE BEST EMBODIMENT OF THE INVENTION

- [0008] Examples of the starting material hydrocarbon include paraffin having secondary carbon, olefin, cycloparaffin and arylalkyl hydrocarbon. Specific preferable examples of the starting material include, but are not limited to, isobutene, etc. as paraffin having secondary carbon; pentene, isobutene, etc. as olefin; cyclopentane, cyclohexane, etc. as cycloparaffin; and cumene, cymene, etc. as arylalkyl hydrocarbon.
- 40

[0009] The compound represented by the following general formula (I) may be cited as an example of the arylalkyl hydrocarbon:

- 45 Compound of the general formula (I):



- 55 (wherein P and Q stand for hydrogen or the alkyl group, which may be the same or different from each other; x stands for a whole number of 1 to 3; and Ar stands for the aromatic hydrocarbon group of the x value.)

[0010] In the general formula (I), it is preferable that at least either of P and Q is the alkyl group, especially preferable that both of them are alkyl groups. Especially the methyl group is preferable as the aforementioned alkyl group. Further,

a hydrocarbon group having the x value that can be derived from benzene, naphthalene, biphenyl, diphenyl ether, etc., preferable a hydrocarbon group of the x value that is derived from benzene or naphthalene, may be cited as examples of the aromatic hydrocarbon group.

[0011] Therefore, in the present invention, preferable examples of the arylalkyl hydrocarbon include, but not limited to, diisopropylbenzenes such as cumene, cymene, m-diisopropylbenzene and p-di-isopropyl benzene, triisopropylbenzenes such as 1,3,5-tri-isopropylbenzene, ethylbenzene, sec-butylbenzene, sec-butylethylbenzene, isopropyl naphthalenes, diisopropyl naphthalene such as 2,6-diisopropyl naphthalene, isopropyl diphenyls, diisopropylbiphenyls such as 4,4'-diisopropylbiphenyl, and mixtures of not less than two of these. Cumene is most preferable.

[0012] The compound of the present invention that can capture radicals refers to a compound having the ability to capture radicals, and it does not matter whether the compound itself possesses such ability or such ability is given to the compound under reaction conditions.

[0013] Examples of the compound of the present invention that can capture radicals include radicals of oxygen, nitrogen, phosphorus, sulfur, carbon and silicon, and the compounds that form these radicals in the reaction system.

[0014] In the preparation of the present invention, either those radicals which are stable at room temperature or those compounds which form radicals under reaction conditions may be used.


[0015] In the reaction of oxidizing hydrocarbon by a gas containing oxygen, various by-products are formed. In the case of the example of the autoxidation of cumene, dimethylphenyl carbinol, acetophenone and dicumylperoxide may be cited as examples of such by-products. It is presumed that there are alkyl radicals such as cumyl radical and cumenylhydroperoxy radical, hydroxy radical, etc. present in the process of reaction. It is presumed in the present invention that the compound that can capture radicals acts on these radicals in some respects, such as capturing the radicals inducing the reaction of the formation of such by-products, and consequently the selectivity of the hydroperoxide corresponding to the raw material hydrocarbon is improved.

[0016] In selecting the compound that can capture radicals in the present invention, it is possible to use the difference in the SOMO (single occupied molecular orbit) energy level ($\Delta \epsilon$ (SOMO)) between the radical of or derived from said compound that can capture radicals, and the radicals to be captured. The SOMO energy level is calculated by the method as described below. It is desirable to select such compound capable of capturing radicals that the difference in the SOMO energy level, $\Delta \epsilon$ (SOMO), is normally 0 to 10eV, preferably 0 to 4eV, more preferably 0 to 1eV.

[0017] For example, it is desirable to select such compound capable of capturing radicals in the present invention that its relationship with the radical (formula 1-2 below) corresponding to the hydrocarbon to be oxidized is as follows:

$$\Delta \epsilon \text{ (SOMO)} = | \epsilon a \text{ (SOMO)} - \epsilon b \text{ (SOMO)} | = 0 \text{ to } 10\text{eV}$$

wherein ϵa (SOMO) stands for the SOMO energy level of the compound cable of capturing radicals or the radical formed therefrom, and ϵb (SOMO) stands for the SOMO energy level of the radical corresponding to the hydrocarbon to be oxidized.



(wherein P and Q, standing for hydrogen or the alkyl group, may be the same or different from each other; x stands for a whole number of 1 to 3; and Ar stands for the aromatic hydrocarbon group of the x value.)

ϵ (SOMO) Calculation Method:

[0018] The structural calculation of a radical is made by a semiempirical molecular orbit method (MNDO-PM3 method: MOPAC program) to calculate ϵ (SOMO).

[0019] The ϵ (SOMO) of those which are not a radical is determined by the structural calculation of the corresponding radical.


[0020] As the aforementioned $\Delta \epsilon$ (SOMO) becomes lower, the hydrocarbon radical is easier to capture. The $\Delta \epsilon$ (SOMO) is normally in the range of 0 to 10eV, preferably 0 to 4eV, more preferably 0 to 1eV.

[0021] The use of a compound having a proper value of $\Delta \epsilon$ (SOMO) has the characteristic that in comparison with the case of adding no such compound, the selectivity goes up over the selectivity of hydroperoxide in other reaction at the same rate of accumulation. Because of this, the use of such compound enables hydrocarbon to be converted at

high selectivity and at a high concentration.

[0022] Oxygen radical and a compound that forms oxygen radical in the reaction system can be cited as one specific example of the compound that can capture radicals.

[0023] As the oxygen radical of the present invention, the compound represented by the general formula (II):



(wherein Z stands for nitrogen, sulfur or phosphorus; X stands for any substitutional group; and n stands for a whole number of 1 to 4 satisfying the value of Z) or a compound that forms the compound represented by the general formula (II) in the reaction system can be cited.

[0024] As the substitutional group X of the general formula (II), the following can be cited:

X, which may be the same or different from each other, may be hydrogen atom, halogen atom, hydrocarbon group, heterocyclic compound residue, oxygen-containing group, nitrogen-containing group, boron-containing group, sulfur-containing group, phosphorus-containing group or silicon-containing group, or a ring formed by not less than two of these being connected to each other.

[0025] As halogen atom, fluorine, chlorine, bromine and iodine.

[0026] Specific examples of the hydrocarbon group include straight-chain or branched alkyl groups having 1 to 30, preferably 1 to 20, carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, neopentyl and n-hexyl; straight-chain or branched alkenyl groups having 2 to 30, preferably 2 to 20, carbon atoms, such as vinyl, aryl and isopropenyl; straight-chain or branched alkynyl group having 2 to 30, preferably 2 to 20 carbon atoms, such as ethynyl and propargyl; cyclic saturated hydrocarbon groups having 3 to 30, preferably 3 to 20, carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and adamantyl; cyclic unsaturated hydrocarbon group having 5 to 30 carbon atoms such as cyclopentadienyl, indenyl and fluorenyl and aryl groups having 6 to 30, preferably 6 to 20, carbon atoms, such as phenyl, benzyl, naphthyl, biphenyl, terphenyl, phenanthryl and anthracenyl.

[0027] Hydrogen atom in the aforementioned hydrocarbon group may be substituted by halogen atom. Examples of such hydrocarbon group include halogenated hydrocarbon groups having 1 to 30, preferably 1 to 20, carbon atoms, such as trifluoromethyl, pentafluorophenyl and chlorophenyl.

[0028] Furthermore, the aforementioned hydrocarbon group may be substituted by other hydrocarbon groups mentioned above. Examples of such hydrocarbon group include aryl-group-substituted alkyl groups such as benzyl and cumyl; alkyl-group-substituted aryl groups such as tolyl, isopropylphenyl, t-butylphenyl, dimethylphenyl and di-t-butylphenyl; and substituted aryl groups in which an aryl group is substituted by alkoxy group, aryl group or aryloxy group. Moreover, the aforementioned hydrocarbon groups may contain the heterocyclic compound residue, oxygen-containing group, nitrogen-containing group, boric acid-containing group, sulfur-containing group, phosphorus-containing group or silicon-containing group mentioned below.

[0029] Out of them, especially preferable hydrocarbon groups are particularly straight-chain or branched alkyl groups having 1 to 30, preferably 1 to 20, carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, neopentyl and n-hexyl; aryl groups having 6 to 30, preferably 6 to 20, carbon atoms, such as phenyl, naphthyl, biphenyl, terphenyl, phenanthryl and anthracenyl; and substituted aryl groups having 1 to 5 of substituents such as alkyl group or alkoxy group having 1 to 30, preferably 1 to 20 carbon atoms, and aryl group or aryloxy group having 6 to 30, preferably 6 to 20 carbon atoms.

[0030] Examples of the heterocyclic compound residue include residues of nitrogen-containing compounds such as pyrrole, pyridine, pyrimidine, quinoline and triazine; oxygen-containing compounds such as furan and pyrane; residues of sulfur-containing compounds such as thiophene; and groups having these heterocyclic compound residues further substituted by substitutional groups such as the alkyl group and alkoxy group having 1 to 30, preferably 1 to 20 carbon atoms.

[0031] Examples of the oxygen-containing group include the alkoxy group, aryloxy group, ester group, acyl group, carboxyl group, carbonate group, hydroxy group, peroxy group and carboxylic acid anhydride group.

[0032] Examples of the alkoxy group, out of these, include the alkoxy group having 1 to 30 carbon atoms, specifically, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy and tert-butoxy. Examples of the aryloxy group include the aryloxy group having 6 to 30 carbon atoms, specifically, phenoxy, 2,6-dimethylphenoxy and 2,4,6-trimethylphenoxy.

Examples of the ester group include the ester group having 1 to 30 carbon atoms, specifically, acetyloxy, benzoyloxy, methoxycarbonyl, phenoxycarbonyl and p-chlorophenoxycarbonyl. Examples of the acyl group include the acyl group having 1 to 30 carbon atoms, specifically, the formyl group, acetyl group, benzoyl group, p-chlorobenzoyl group and p-methoxybenzoyl group.

[0033] Examples of the nitrogen-containing group include the amino group, imino group, amide group, imide group, hydrazino group, hydrazono group, nitro group, nitroso group, cyano group, isocyano group, ester cyanate group, amidino group, diazo group and amino group having turned into ammonium salt.

5 [0034] Examples of the amino group, out of these, include the amino group having 0 to 30 carbon atoms, specifically, dimethylamino, ethylmethylamino and diphenylamino. Examples of the imino group include the imino group having 1 to 30 carbon atoms, specifically, methylimino, ethylimino, propylimino, butylimino and phenylimino. Examples of the amido group include the amido group having 1 to 30 carbon atoms, specifically, acetoamido, N-methylacetoamido and N-methylbenzamino. Examples of the imido group is the imido group having 2 to 30 carbon atoms, specifically, acetoimido and benzimido.

10 [0035] Examples of the boron-containing group include the boranediol group, boranetolyl group and diboranyl group.

[0036] Examples of the sulfur-containing group include the mercapto group, thioester group, dithioester group, alkylthio group, arylthio group, thioacyl group, thioether group, thiocyanate group, isocyanate group, sulfonester group, sulfonamide group, thiocarboxyl group, dithiocarboxyl group, sulfo group, sulfonyl group, sulfinyl group and sulphenyl group.

15 [0037] Examples of the thioester group, out of these, include the thioester group having 1 to 30 carbon atoms, specifically, acetylthio, benzoylthio, methylthiocarbonyl and phenylthio-carbonyl. Examples of the alkylthio group include the alkylthio group having 1 to 30 carbon atoms, specifically, methylthio and ethylthio. Examples of the sulfonamide group include the sulfonamide group having 0 to 30 carbon atoms, specifically, phenylsulfonamide, N-methylsulfonamide and N-methyl-p-toluene-sulfonamide. Examples of the arylthio group include the arylthio group having 6 to 30 carbon atoms, specifically, phenylthio, methylphenylthio and naphthylthio. Examples of the sulfonester group include the sulfonester group having 1 to 30 carbon atoms, specifically, methyl sulfonic acid, ethyl sulfonic acid and phenyl sulfonic acid.

[0038] Examples of the phosphorus-containing group include the phosphide group, phosphoryl group, thiophosphoryl and phosphato group.

25 [0039] Examples of the silicon-containing group include the silyl group, siloxy group, hydrocarbon-substituted silyl group, hydrocarbon-substituted siloxy group, specifically, methyl silyl, dimethylsilyl, trimethylsilyl, ethylsilyl, diethylsilyl, triethylsilyl, diphenylmethylsilyl, triphenylsilyl, dimethylphenylsilyl, dimethyl-t-butylsilyl, and dimethyl(pentafluorophenyl)silyl. Out of these, methyl silyl, dimethylsilyl, trimethylsilyl, ethylsilyl, diethylsilyl, triethylsilyl, dimethylphenylsilyl and triphenylsilyl are preferable. Especially, trimethylsilyl, triethylsilyl, triphenylsilyl and dimethylphenylsilyl are preferable.

30 Specific examples of the hydrocarbon-substituted siloxy group include trimethylsiloxy.

[0040] Examples of the general formula (II) include, but are not limited to, the following, for example:


35

40


45

50


55



[0041] In the general formula (II), a compound represented by the following general formula (III) may be cited as an example of nitroxide radical.



(wherein m stands for a whole number of 0 to 3; and -D- and -D'- stand each separately for:



A, B, X¹, X², R¹, R², R³, R⁴ and W stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, Phosphorus, silicon and halogen; and each ring member carbon atom or each ring member nitrogen atom may form a double bond between adjoining atoms).

[0042] The substitutional group in the general formula (III) may be the same group as cited as an example of the substitutional group X in the aforementioned general formula (II). Preferable examples of the hydrogen or the substitutional groups of the general formula (III) are hydrogen, alkyl group, aryl group, halogen, cyano group, amino group, isothiocyanic acid group, -COORa (wherein Ra stands for hydrogen, alkyl group and aryl group), (di)alkylaminoalkyl group, hydroxyl group, hydroxyalkyl group, alkoxy group, arylalkoxy group, -CONRbRc (wherein Rb and Rc are each separately hydrogen, alkyl group and aryl group), oxo group (=O), maleicimide group, phosphoric acid group, =NH group or bivalent group.

[0043] The examples of the aforementioned nitroxide radicals may include a compound having a structure in which two or more structures of the general formula (III) are crosslinked through any of the groups of X¹, X², A, B and W.

[0044] In the general formula (III), it is preferable that all of R¹, R², R³ and R⁴ are groups selected from alkyl groups. For example, 2,2,6,6-teraalkylpiperidinoxy radical, 4-hydroxy-2,2,6,6-tetraalkyl piperidinyloxy radical, bis-(2,2,6,6-teraalkyl piperidinoxy)-sebacate or 2,2,5,5-teraalkyl pyrrolidinoxy radical or the said compound substituted by the aforementioned group may be cited as preferable compounds.

[0045] Specific examples of the compound of the general formula (III) include, but are not limited to, the following:

5

10

15

20

25

30


35

40

45

50

55



5

10

15

20

25

30


35


40

45

50


55





(wherein Me stands for the methyl group; Et stands for the ethyl group; and Ph stands for the phenyl group: This hereinafter applies to the present invention. Further, p and q stand for a whole number of 0 to 30, respectively.)

[0046] As the compound that turns into nitroxide radical under reaction conditions, the compound that turns into nitroxide radical represented by the aforementioned general formula (III) under reaction conditions or the compounds represented by the following general formulas (V), (VI) and (VII).




(wherein n stands for a whole number of 0 to 4; and R_5 and R^6 stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen.)

[0047] Each of the substitutional groups in the general formula (V) may be the same group as cited as an example of the substitutional group X in the aforementioned general formula (II). A preferable example of R_5 may be halogen, alkyl group, arylalkyl group, alkoxy group or arylalkoxy group, and a preferable example of R^6 may be hydrogen or the alkyl group.

[0048] As examples of each of substitutional groups of the general formula (V), halogen includes F, Cl, Br and I; the alkyl group preferably includes the alkyl group having 1 to 10 carbon atoms, such as the methyl group, ethyl group, n-propyl group, i-propyl group, n-butyl group, i-butyl group, sec-butyl group, t-butyl group and isoamyl group; the arylalkyl group includes cumyl group; the alkoxy group preferably includes the alkoxy group having 1 to 10 carbon atoms, such as methoxy group, ethoxy group, n-propoxy group, i-propoxy group, n-butoxy group, i-butoxy group, sec-butoxy group and t-butoxy group; and the arylalkoxy group includes cumyloxy group.


[0049] Specific examples of the compound of the general formula (V) include, but are not limited to, N-hydroxyphthalimide and N-methoxyphthalimide:



(wherein R^7 , R^8 and R^9 stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen.)

[0050] Each of the substitutional groups in the general formula (VI) may be the same group as cited as an example of the substitutional group X in the aforementioned general formula (II). Preferable examples of the substitutional groups are halogen, alkyl group, or haloalkyl group for R^7 , R^8 and R^9 .

[0051] As examples of each of substitutional groups of the general formula (VI), halogen includes F, Cl, Br and I; the alkyl group preferably includes the alkyl group having 1 to 10 carbon atoms, such as the methyl group, ethyl group, n-propyl group, i-propyl group, n-butyl group, i-butyl group, sec-butyl group, t-butyl group and isoamyl group; and the haloalkyl group includes trifluoromethyl group.







10 (wherein l stands for a whole number of 0 to 3; and Y, R¹⁰ and R¹¹ stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen.)

15 [0052] Each of the substitutional groups in the general formula (VII) may be the same group as cited as an example of the substitutional group X in the aforementioned general formula (II). Preferable examples of the substitutional groups are halogen, alkyl group, arylalkyl group, alkoxy group or arylalkoxy group.

20 [0053] Halogen includes F, Cl, Br and I; the alkyl group preferably includes the alkyl group having 1 to 30 carbon atoms, such as the methyl group, ethyl group, n-propyl group, i-propyl group, n-butyl group, i-butyl group, sec-butyl group, t-butyl group and isoamyl group; the arylalkyl group includes cumyl group; the alkoxy group preferably includes the alkoxy group having 1 to 30 carbon atoms, such as methoxy group, ethoxy group, n-propoxy group, i-propoxy group, n-butoxy group, i-butoxy group, sec-butoxy group and t-butoxy group; and the arylalkoxy group includes cumyloxy group.



[0054] Specific examples of the compound that can capture radicals include a nitrogen radical or a compound that forms a nitrogen radical in the reaction system.

25 [0055] Specific examples of the compound that can capture these radicals include, but are not limited to, the following:




45 [0056] Specific examples of the compound that can capture radicals include a phosphorus radical or a compound that forms a phosphorus radical in the reaction system.

[0057] Specific examples of the compound that can capture these radicals include, but are not limited to, the following:




[0058] Specific examples of the compound that can capture radicals include a sulfur radical or a compound that forms a sulfur radical in the reaction system.

[0059] Specific examples of the sulfur radical include, but are not limited to, the following:






[0060] Specific examples of the compound that can capture radicals include a carbon radical or a compound that forms a carbon radical in the reaction system.

[0061] Specific examples of the carbon radical include, but are not limited to, the following:




[0062] Specific examples of the compound that can capture radicals include a silicon radical or a compound that forms a silicon radical in the reaction system.


[0063] Specific examples of the silicon radical include, but are not limited to, the following:



[0064] The compound that can capture radicals include secondary amine. Preferably the secondary amine represented by the following general formula (IV) may be cited.



(wherein m stands for a whole number of 0 to 3; -D- and -D'- stand each separately:



A, B, X¹, X², R¹, R², R³, R⁴ and W stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen; and each ring member carbon atom or each ring member nitrogen atom may form a double bond between adjoining atoms).









[0065] The substitutional group in the general formula (IV) may be the same group as cited as an example of the substitutional group X in the aforementioned general formula (II). Preferable examples of the hydrogen or other substitutional groups of the general formula (IV) are hydrogen, alkyl group, aryl group, halogen, cyano group, amino group, isothiocyanic acid group, -COORa (wherein Ra stands for hydrogen, alkyl group and aryl group), (di)alkylaminoalkyl group, hydroxyl group, hydroxyalkyl group, alkoxy group, arylalkoxy group, -CONRbRc (wherein Rb and Rc are each separately hydrogen, alkyl group and aryl group), oxo group (=O), maleimide group, phosphoric acid group, =NH group or bivalent group.

[0066] The substitutional groups, which may be the same or different from each other, may be hydrogen atom, halogen atom, hydrocarbon group, heterocyclic compound residue, oxygen-containing group, nitrogen-containing group, boron-containing group, sulfur-containing group, phosphorus-containing group or silicon-containing group, or a ring formed by not less than two of these being connected to each other.

[0067] The examples of the aforementioned secondary amine may include a compound having a structure in which two or more structures of the general formula (IV) are crosslinked through any of the radicals of X¹, X², A, B and W.

[0068] In the general formula (IV), the preferable compounds are that all of R¹, R², R³ and R⁴ are groups selected from the alkyl group. For example, they include 2,2,6,6-tetraalkyl piperidine, 2,2,6,6-tetraalkyl-4-piperidinol, bis(2,2,6,6-tetraalkyl piperidinyl)-sebacate or 2,2,5,5-tetraalkyl pyrrolidine, or the said compound substituted by the aforementioned group.

[0069] Specific examples of the secondary amine include, but are not limited to, the following:



5

10

15

20

25

30


35


40

45


50

55





[0070] For the compound that can capture radicals, hindered phenols represented by the general formula (VIII) may be cited.



10 (wherein p stands for a whole number of 0 to 3; R¹² and R¹³ stand each separately for a substitutional group; and z stands for hydrogen or another substitutional group.)

15 [0071] It is preferable that R¹² and R¹³ are alkyl group having 1 to 30 carbon atoms or the aryl group having 1 to 30 carbon atoms, and Z is hydrogen, halogen, alkyl group or aryl group.

[0072] The substitutional group in the general formula (VIII) may be the same group as an example of the substitutional group X in the aforementioned general formula (II).

[0073] Specific examples of the compound of the general formula (VIII) include, but are not limited to, the following:

20

25

30


35

40


45

50

55



[0074] As the compound that can capture radicals, phosphine oxides may be cited. Phosphine oxides of the present invention include the compounds of the general formula (IX)
The compounds of the general formula (IX)




(wherein R^{14} , R^{15} and R^{16} stand each separately for hydrogen or another substitutional group.)

[0075] It is preferable desirable that R^{14} , R^{15} and R^{16} are the alkyl group or the aryl group.

[0076] The substitutional group in the general formula (IX) may be the same group as cited as an example of the substitutional group X in the aforementioned general formula (II).


[0077] Specific examples of the compound of the general formula (IX) include, but are not limited to, the following:



[0078] Furthermore, in the process for preparation of the present invention, either phosphineoxide which is stable at room temperature or a compound which turns into phosphineoxide under reaction conditions may be used.

[0079] As the compound that turns into phosphineoxide under reaction conditions, the compound represented by the


general formula (X) may be cited.



(wherein R^{17} , R^{18} and R^{19} stand each separately for hydrogen or another substitutional group.)


[0080] The substitutional group in the general formula (X) may be the same group as an example of the substitutional group X in the aforementioned general formula (II). It is preferable that these substitutional groups are the alkyl group or the aryl group.

[0081] Specific examples of the compound of the general formula (X) include, but are not limited to, triphenylphosphine and tributylphosphine.



[0082] A spin trap agent may be used as the compound that can capture radicals. The spin trap agent is a compound which is used for identifying a radical with a short life that cannot be observed directly by means of ESR, and forms a spin trap compound radical observable with the ESR with the radical to enable to identify the original radical with short life time by ESR analysis.


[0083] Specific examples of the compound include, but are not limited to, nitron and nitroso compound as shown below.



[0084] In this invention, the oxidation of hydrocarbon is conducted in the presence of at least one compound that can capture radicals. Two or more compounds that can capture radicals can be used.

[0085] Supposing the compound that can capture radicals has captured a radical, the enthalpy (ΔH) to break the bond between the compound and the radical can be determined by calculation by the method as described below. The appropriate range of ΔH depends on the type of the radical to be captured, and the value may be used to select the compound that can capture radicals of the present invention.

[0086] Satisfactory results can be obtained by selecting as the compound of the present invention that can capture radicals such compound that the value of enthalpy (ΔH) is -30 to 50KJ/mol when calculated on the basis of the assumption that the radical to be captured is the radical corresponding to the hydrocarbon to be oxidized (formula I-2). The value of the enthalpy (ΔH) is preferably -25 to 50KJ/mol, more preferably -20 to 50KJ/mol. Using the compound that can capture those radicals whose ΔH is within such appropriate range produces satisfactory results as intended by the present invention.



H (heat of formation) calculation method:

[0087] H was calculated by conducting structural optimization for the structure showing the smallest amount of energy (most stable structure) by semiempirical molecular orbital method(AM1 method: MOPAC program) after carrying out conformation search by the molecular force field method (CHARMm force field: QUANTA program).

[0088] The selection of the compound that can capture radicals has the figures within the appropriate range of $\Delta \epsilon$ (SOMO) aforementioned and the value of the enthalpy (ΔH) is most preferable in this invention.

[0089] Next, given below is an explanation of the reaction conditions of the present invention.

[0090] The usage of the compound that can capture radicals or the compound that turns into the compound that can capture radicals in the reaction system in the present invention is normally in the range of 0.00001 to 5.0 parts by weight, preferably in the range of 0.0001 to 0.1 parts by weight, per 100 parts by weight of the starting material hydrocarbon.

[0091] As the oxygen-containing gas as the oxidizing agent, air is normally used, but oxygen or any mixed gas of oxygen and nitrogen may also be used.

[0092] The reaction may normally be conducted at atmospheric pressure, but it may also be conducted under elevated pressure. The reaction temperature is normally in the range of 40 to 120°C, preferably in the range of 50 to 100°C.

[0093] Furthermore, this reaction may be conducted in the presence of a basic compound in the state of a solid or an aqueous solution. Examples of the basic compounds include sodium carbonate, sodium hydrogencarbonate, sodium hydroxide, potassium carbonate, potassium hydrogencarbonate, potassium hydroxide, barium oxide and magnesium oxide. The amount of the basic compound used in the reaction is normally 0.0001 to 10.0 parts by weight, preferably 0.001 to 5.0 parts by weight, per 100 parts by weight of the starting material hydrocarbon.

[0094] The reaction in the present invention may be conducted by either batch or continuous method. In the case of conducting the reaction by the batch method, if the starting material hydrocarbon is a liquid at the reaction temperature, the aforementioned compound that can capture radicals or the aforementioned compound that turns into the compound that can capture radicals in the reaction system is added to the starting material and normally made to undergo oxida-

tion reaction by blowing air into the material with heating and agitation. If required, an inactive organic solvent may be used as the reaction solvent in the reaction. On the other hand, if the starting material arylalkyl hydrocarbon is a solid at the reaction temperature, it is made to be dissolved in an inactive organic solvent to form a solution, and the aforementioned compound that can capture radicals or the aforementioned compound that turns into the compound that can capture radicals in the reaction system may be added to the solution and made to undergo oxidation reaction by blowing air into the material with heating and agitation.

[0095] Further, in the present invention, a small amount of hydroperoxide which corresponds to the hydrocarbon may be present, as required, as the initiator in the hydrocarbon at the time of the start of the reaction.

[0096] A catalyst such as a transition metal complex may also be used additionally as required.

[0097] Such catalyst may be formed as a fixed bed, and the starting material hydrocarbon or its solution and air may be caused to pass through the fixed bed, while being mixed.

[0098] According to the method of the present invention, the formed hydroperoxide can be easily recovered from the reaction mixture by distillation or any other conventional means after the completion of the reaction and after filtering out the catalyst as required.

Examples

[0099] Given below is a specific description of examples of preparing organic hydroperoxides from arylalkyl hydrocarbon. However, the present invention is not limited in any way by the following examples:

Example 1

[0100] 18 mg of 2,2,6,6-tetramethylpiperidinoxyl were added to a mixture of 144 g of cumene and 36 g of cumenehydroperoxide, and all of them were dissolved. 90 g of 0.05 wt% sodium carbonate were added to it, and this mixture was heated under 6 K pressure to 105°C. Then air was blown into it at a rate of 400 ml/min with vigorous agitation for 3 hours to oxidize cumene.

[0101] The rate of accumulation of cumenehydroperoxide during 3 hours from the start of the reaction was determined by iodometry and gas chromatography to be 5.0 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 91 mol%.

[0102] The $\Delta \epsilon$ and enthalpy (ΔH) of the 2,2,6,6-tetramethylpiperidinoxyl used and the cumyl radicals corresponding to cumene were calculated to be 0.2eV and -6.4KJ/mol, respectively.

Example 2

[0103] The reaction was conducted in the same manner as Example 1 except that di-*t*-butylnitroxide was used in place of 2,2,6,6-tetramethylpiperidinoxyl.

[0104] The rate of accumulation of cumenehydroperoxide during 3 hours from the start of the reaction was determined by iodometry and gas chromatography to be 5.2 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 90 mol%.

[0105] The enthalpy (ΔH) of the di-*t*-butylnitroxide used and the cumyl radicals corresponding to cumene were calculated to be -15.0KJ/mol.

Example 3

[0106] The reaction was conducted in the same manner as Example 1 except that N-hydroxyphthalimide was used in place of 2,2,6,6-tetramethylpiperidinoxyl.

[0107] The rate of accumulation of hydroperoxide during 3 hours from the start of the reaction was determined by iodometry and gas chromatography to be 5.0 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 87 mol%.

Example 4

[0108] The reaction was conducted in the same manner as Example 1 except that potassium nitrosodisulfonate was used in place of 2,2,6,6-tetramethylpiperidinoxyl.

[0109] The rate of accumulation of cumenehydroperoxide during 3 hours from the start of the reaction was determined by iodometry and gas chromatography to be 5.1 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 89 mol%.

Example 5

[0110] 50 mg of 2,2,6,6-tetramethyl-4-piperidinol were added to a mixture of 126 g of cumene and 54 g of cumenehydroperoxide, and all of them were dissolved. 5.6 g of purified water were added to it, and this mixture was heated under 6 K pressure to 105°C. The cumene was oxidized by continuous reaction at a residence time of 1 hour while blowing air into it at a rate of 400 ml/min with vigorous agitation.

[0111] The rate of accumulation of cumenehydroperoxide after 4, 5 and 6 hours from the start of the reaction was determined and averaged by iodometry and gas chromatography to be 3.7 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined and averaged by high performance liquid chromatography to be 92 mol%.

Example 6

[0112] The reaction was conducted in the same manner as Example 5 except that 36 mg of bis-(2,2,6,6-tetramethyl-4-piperidinyl)-sebacate was added in place of 2,2,6,6-tetra-methyl-4-piperidinol.

[0113] The rate of accumulation of cumenehydroperoxide after 4, 5, and 6 hours from the start of the reaction was determined and averaged by iodometry and gas chromatography to be 3.8 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined and averaged by high performance liquid chromatography to be 90 mol%.

Example 7

[0114] 5 g of pure water and 36 mg of 2, 4, 6-triphenylphenol were added to a mixture of 122 g of cumene and 53 g of cumenehydroperoxide, and all of them were dissolved as an oil phase. This mixture was heated under 6 K pressure to 105°C. The cumene was oxidized while blowing air into it at a rate of 180 ml/min with vigorous agitation.

[0115] A solution of 36 mg of 2, 4, 6-triphenylphenol dissolved in a mixture of 122 g of cumene and 53 g of cumenehydroperoxide, and pure water were added into the reactor at rates of 175 g/hour and 5 g/hour, respectively, in such manner that the residence time is 1 hour of continuous reaction. The reaction liquid was collected by overflow into another vessel three times every hour after 4 hours from the start of the reaction.

[0116] The rate of accumulation of cumenehydroperoxide was determined by iodometry and gas chromatography. The average of three determination values was 3.8 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography. The average of three determination values was 89 mol%.

[0117] The $\Delta \varepsilon$ (SOMO) and enthalpy (ΔH) between the 2,4,6-triphenylphenoxy radical presumed to be formed from the 2, 4, 6-triphenylphenol used and cumyl radical corresponding to cumene were calculated to be 0.4eV and 16.5KJ/mol, respectively.

Example 8

[0118] The reaction was conducted in the same manner as Example 7 except that 2, 6-di-t-butyl-4-methylphenol was added in place of 2, 4, 6-triphenylphenol.

[0119] The rate of accumulation of cumenehydroperoxide was determined by iodometry and gas chromatography. The average of three determination values was 3.5 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 90 mol%.

Example 9

[0120] 5 g of pure water and 36 mg of triphenylphosphineoxide were added to a mixture of 122 g of cumene and 53 g of cumenehydroperoxide, and all of them were dissolved as an oil phase. This mixture was heated under 6 K pressure to 105°C. Then air was blown into it at a rate of 180 ml/min with vigorous agitation to oxidize cumene.

[0121] A solution of 36 mg of triphenylphosphineoxide dissolved in a mixture of 122 g of cumene and 53 g of cumenehydroperoxide and pure water were added to the reactor at rates of 175 g/hour and 5 g/hour, respectively, in such manner that the residence time is 1 hour of continuous reaction. The reaction liquid was collected by overflow into another vessel three times every hour after 4 hours from the start of the reaction.

[0122] The rate of accumulation of cumenehydroperoxide was determined by iodometry and gas chromatography. The average of three determination values was 4.5 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography. The average of three determination values was 90 mol%.

Example 10

[0123] The reaction was conducted in the same manner as described in Example 9 except that tri-*t*-butylphosphine-oxide was used as an additive in place of triphenylphosphineoxide.

[0124] The rate of accumulation of cumenehydroperoxide was determined by iodometry and gas chromatography. The average of three determination values was 3.5 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 90 mol%.

Comparative Example 1

[0125] The reaction was conducted in the same manner as Example 1 except that 2, 2, 6, 6-tetramethylpiperidinoxy was not used.

[0126] The rate of accumulation of cumenehydroperoxide during 3 hours from the start of the reaction was determined by iodometry and gas chromatography to be 5.5 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 84 mol%.

Comparative Example 2

[0127] The reaction was conducted in the same manner as Comparative Example 1 except that the reaction temperature was set at 100°C.

[0128] The rate of accumulation of cumenehydroperoxide during 3 hours from the start of the reaction was determined by iodometry and gas chromatography to be 4.1 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 88 mol%.

Comparative Example 3

[0129] The reaction was conducted in the same manner as Example 5 except that 2, 2, 6, 6-tetramethyl-4-piperidinol was not used.

[0130] The rate of accumulation of cumenehydroperoxide during 3 hours from the start of the reaction was determined and averaged by iodometry and gas chromatography to be 5.5 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined and averaged by high performance liquid chromatography to be 86 mol%.

Comparative Example 4

[0131] The reaction was conducted in the same manner as Comparative Example 3 except that the reaction temperature was set at 100°C.

[0132] The rate of accumulation of cumenehydroperoxide during 4, 5 and 6 hours from the start of the reaction was determined and averaged by iodometry and gas chromatography to be 3.9 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined and averaged by high performance liquid chromatography to be 89 mol%.

Comparative Example 5

[0133] The reaction was conducted in the same manner as Example 7 except that 2, 4, 6-triphenylphenol was not used.

[0134] The rate of accumulation of cumenehydroperoxide was determined by iodometry and gas chromatography. The average of three determination values was 5.5 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 86 mol%.

Comparative Example 6


[0135] The reaction was conducted in the same manner as Comparative Example 5 except that the reaction temperature was set at 100°C.

[0136] The rate of accumulation of cumenehydroperoxide was determined by iodometry and gas chromatography. The average of three determination values was 3.9 wt%/hr. Furthermore, the selectivity of the cumenehydroperoxide which was formed in the reaction was determined by high performance liquid chromatography to be 88 mol%.

Claims

1. A process for preparing hydroperoxides, which comprises oxidizing hydrocarbon by a gas containing oxygen in the presence of at least one compound that can capture radicals and converting it selectively to corresponding hydroperoxides.

2. The process as defined in Claim 1, wherein arylalkyl hydrocarbon represented by the general formula (I)



(wherein P and Q, standing for hydrogen or the alkyl group, may be the same or different from each other; x stands for a whole number of 1 to 3; and Ar stands for the aromatic hydrocarbon group of the x value) is oxidized by a gas containing oxygen and converting it selectively to corresponding arylalkylhydroperoxides.

3. The process as defined in Claim 2, wherein the arylalkyl hydrocarbon represented by the general formula (I) is cumene, cymene, m-diisopropylbenzene, p-diisopropylbenzene, 1,3,5-triisopropylbenzene, isopropyl naphthalene, diisopropyl naphthalene, isopropylbiphenyl, diisopropylbiphenyl or a mixture of two or more of these.

4. The process as defined in Claim 3, wherein arylalkyl hydrocarbon is cumene.

5. The process as defined in any claim of Claim 1 to 4, wherein the compound that can capture radicals is radicals of oxygen, nitrogen, phosphorus, sulfur, carbon or silicon or a compound that forms radicals of these in the reaction system.

6. The process as defined in Claim 5, wherein the compound that can capture radicals is an oxygen radical or a compound that forms an oxygen radical in the reaction system.

7. The process as defined in Claim 5, wherein the compound that can capture radicals is a nitrogen radical or a compound that forms a nitrogen radical in the reaction system.

8. The process as defined in Claim 5, wherein the compound that can capture radicals is a phosphorus radical or a compound that forms a phosphorus radical in the reaction system.


9. The process as defined in Claim 5, wherein the compound that can capture radicals is phosphine oxide or a compound that forms phosphine oxide in the reaction system.

10. The process as defined in Claim 5, wherein the compound that can capture radicals is a sulfur radical or a compound that forms a sulfur radical in the reaction system.

11. The process as defined in Claim 5, wherein the compound that can capture radicals is a carbon radical or a compound that forms a carbon radical in the reaction system.


12. The process as defined in Claim 5, wherein the compound that can capture radicals is a silicon radical or a compound that forms a silicon radical in the reaction system.

13. The process as defined in Claim 6, wherein the compound that can capture radicals is a compound represented by the general formula (II):




(wherein Z stands for nitrogen, sulfur or phosphorus; X stands for any substitutional group; and n stands for a whole number of 1 to 4 satisfying the value of Z) or a compound that forms the compound represented by the general formula (II) in the reaction system.

14. The process as defined in Claim 13, wherein the compound that can capture radicals is a compound represented by the general formula (III):



(wherein m stands for a whole number of 0 to 3; and -D- and -D'- stand each separately for:



A, B, X¹, X², R¹, R², R³, R⁴ and W stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen; and each ring member carbon atom or each ring member nitrogen atom may form a double bond between adjoining atoms).

15. The process as defined in Claim 14, wherein the compound that can capture radicals is a compound having a structure in which two or more structures of the general formula (III) are crosslinked through any of the groups of X¹, X², A and B.

16. The process as defined in Claim 15, wherein the hydrogen or other substitutional groups of the general formula (III) are hydrogen, alkyl group, aryl group, halogen, cyanogen group, amino group, isothiocyanic acid group, -COOR_a (wherein R_a stands for hydrogen, alkyl group and aryl group), (di)alkylaminoalkyl group, hydroxyl group, hydroxyalkyl group, alkoxy group, arylalkoxy group, -CONR_bR_c (wherein R_b and R_c are each separately hydrogen, alkyl group and aryl group), oxo group (= O), maleimide group, phosphoric acid group, =NH group or bivalent group.


17. The process as defined in Claim 14, wherein R¹, R², R³ and R⁴ are the alkyl group in the general formula (III).

18. The process as defined in Claim 17, wherein the compound represented by the general formula (III) is 2,2,6,6-tetraalkyl piperidinoxy radical, 4-hydroxy-2,2,6,6-tetraalkyl piperidinyloxy radical, bis-(2,2,6,6-tetraalkyl piperidinoxy)sebacate or 2,2,5,5-tetraalkyl pyrrolidinoxy radical.


19. The process as defined in Claim 1, wherein the compound that capture radicals is hindered phenol.

20. The process as defined in Claim 1, wherein the compound that capture radicals is secondary amine.

21. The process as defined in Claim 20, wherein secondary amine is a compound represented by the general formula (IV):



(wherein m stands for a whole number of 0 to 3; -D- and -D'- stand each separately:



A, B, X1, X2, R1, R2, R3, R4 and W stand each separately hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen; and each ring member carbon atom or each ring member nitrogen atom may form a double bond between adjoining atoms).

22. The process as defined in Claim 21, wherein secondary amine is a compound having a structure in which two or more structures of the general formula (IV) are crosslinked through any of the radicals of X1, X2, A and B.

23. The process as defined in Claim 21, wherein the hydrogen or other substitutional groups of the general formula (IV) are hydrogen, alkyl group, aryl group, halogen, cyanogen group, amino group, isothiocyanic acid group, -COORa (wherein Ra stands for hydrogen, alkyl group or aryl group), (di)alkylaminoalkyl group, hydroxyl group, hydroxyalkyl group, alkoxy group, arylalkoxy group, -CONRbRc (wherein Rb and Rc are each separately hydrogen, alkyl group or aryl group), oxo group (= O), maleimide group, phosphoric acid group, =NH group or bivalent group.


24. The process as defined in Claims 21 22 or 23, wherein R1, R2, R3 and R4 are the alkyl group in the general formula (IV).

25. The process as defined in Claim 24, wherein the compound represented by the general formula (IV) is 2,2,6,6-teraaalkylpiperidine, 2,2,6,6-tetraalkyl-4-piperidinol, bis-(2,2,6,6-teraaalkylpiperidinyl)-sebacate or 2,2,5,5-teraaalkylpyrrolidine.

26. The process as defined in Claim 1, wherein the SOMO energy levels of the compound that can capture radicals and the radicals to be captured are determined by calculation by the MNDO-PM3 method and the difference between them is used in the selection of the compound that can capture radicals.


27. The process as defined in Claim 26, wherein the difference in the SOMO energy level ($\Delta \epsilon$ (SOMO)) between the compound that can capture radicals and the radicals to be captured is not more than 10eV as calculated by the MNDO-PM3 method.

28. The process as defined in Claim 1, wherein the difference in the SOMO energy level ($\Delta \epsilon$ (SOMO)) between the compound that can capture radicals and the radicals corresponding to hydrocarbons to be oxidized represented by the following formula (1-2) is not less than 0eV and not more than 10eV as calculated by the MNDO-PM3 method.




(wherein P and Q, standing for hydrogen or the alkyl group, may be the same or different from each other; x stands for a whole number of 1 to 3; and Ar stands for the aromatic hydrocarbon group of the x value.)

29. The process as defined in Claim 1, 26, 27 or 28, wherein a compound that the calculated enthalpy (ΔH) value to break the bond between the compound and a radical of the following formula corresponding to the hydrocarbon to be oxidized when assuming the compound has captured the radical, to form the compound and the radical as before the capture, is -30 to 50 KJ/mol, are used as the compound that can capture radicals:




(wherein P and Q, standing for hydrogen or the alkyl group, may be the same or different from each other; x stands for a whole number of 1 to 3; and Ar stands for the aromatic hydrocarbon group of the x value).

30. A process for preparing hydroperoxides by oxidizing arylalkyl hydrocarbon represented by the general formula (I) by a gas containing oxygen in the presence of at least one nitroxide radical, and converting it selectively to corresponding arylalkyl hydroperoxides.




(wherein P and Q, standing for hydrogen or the alkyl group, may be the same or different from each other; x stands for a whole number of 1 to 3; and Ar stands for the aromatic hydrocarbon group of the x value)

31. The process as defined in Claim 30, wherein the nitroxide radical is a compound represented by the general formula (III):




15 (wherein m stands for a whole number of 0 to 3; and -D- and -D'- stand each separately for:




25 A, B, X1, X2, R1, R2, R3, R4 and W stand each separately for hydrogen or a substitutional group comprising an element selected from hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen; and each ring member carbon atom or each ring member nitrogen atom may form a double bond between adjoining atoms).

32. The process as defined in Claim 30, wherein a compound that turns into nitroxide radical in the reaction system is present in the system in place of nitroxide radical.


30 33. The process as defined in Claim 32, wherein a compound that turns into nitroxide radical is at least one compound selected from the compounds of the general formulas (V), (VI) and (VII):



40 (wherein n stands for a whole number of 0 to 4; and R5 and R6 stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen.)



55 (wherein R7, R8 and R9 stand each separately for hydrogen or a substitutional group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen.)



10 (wherein 1 stands for a whole number of 0 to 3; and Y, R10 and R11 stand each separately for hydrogen or a substituent group comprising an element selected from among hydrogen, carbon, oxygen, nitrogen, sulfur, phosphorus, silicon and halogen.)

15 34. The process as defined in claim 30, wherein the hydrocarbon is cumene.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/02935

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl. ⁶ C07C409/10, 409/00, 407/00 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl. ⁶ 409/00-44, 407/00 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP, 48-49735, A (Mitsui Petrochemical Industries, Ltd.), July 13, 1973 (13. 07. 73) (Family: none)	1-34
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search July 27, 1998 (27. 07. 98)		Date of mailing of the international search report August 4, 1998 (04. 08. 98)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)